

A SOCIAL HEALTH ATLAS OF AUSTRALIA

Second Edition

Volume 5: South Australia

John Glover and Sarah Tennant

December 1999

 **Public Health Information Development Unit**

Copyright

© Commonwealth of Australia 1999

This work may be reproduced and used subject to acknowledgement of the source of any material so rep

National Library of Australia Cataloguing in Publication entry

Glover, John, 1945-

A social health atlas of Australia

2nd ed.

Bibliography.

Includes index.

ISBN 0 7308 9021 X (set).

ISBN 0 7308 9026 0 (v. 5).

1. Medical geography – Australia - Maps. 2. Medical - geography – Australia - Statistics. 3. Public health – Australia - Maps. 4. Public health – Australia - Statistics. 5. Health facilities – Australia – Utilization - Maps. 6. Health facilities – Australia – Utilization – Statistics. 7. Health surveys – Australia. I. Tennant, Sarah. II. University of Adelaide. Public Health Information Development Unit. III. Commonwealth Department of Health and Aged Care (Australia). IV. Title.

362.10994

A Social Health Atlas of Australia was produced by the Public Health Information Development Unit, University of Adelaide, South Australia. The project was funded by the Commonwealth Department of Health and Aged Care and supported by the South Australian Department of Human Services.

The majority of data mapped in the atlas, and the maps, was produced by Prometheus Information Pty Ltd, using the HealthWIZ software package. Further details of these and other contributors to the project are noted in the *Acknowledgements*.

Cover design by Julie Johninke using a photograph by Paul Doherty (photograph entitled *Merti Merti Sandhills, Strzelecki Track*).

Printed in Australia by Openbook Publishers.

Related publications and software products

A Social Health Atlas of Australia, 1992, Vols 1 & 2

HealthWIZ: details available at www.prometheus.com.au

Social health atlas World Wide Web site: www.publichealth.gov.au

Foreword

The publication of this second edition of ***A Social Health Atlas of Australia*** brings together a wide range of information about the health status of Australians by region, and the health service use by the Australian population.

By presenting the data as maps, the atlas provides a graphical image of the distribution of health status, and differences in the patterns and levels of access to and use of health services at the local level throughout the cities, towns, and rural and remote areas of Australia. The format of the atlas makes the information easy to understand and readily accessible to a broad group of users, including public health planners, providers, researchers, students and the general public.

The graphs of the newly developed Accessibility/Remoteness Index for Australia (ARIA) provide useful information for communities, as well as practitioners and managers in the health sector, to better understand the differences in the statistics that describe health status and health service use.

This data is essential for policy development and local area planning, and for monitoring and evaluating health services. It is also of major importance for resource allocation at the broadest level, and between areas, services and population groups. The maps and tabulations presented in this atlas represent a major compilation of information for these purposes.

I congratulate all those who have contributed to this important project.



Dr Michael Wooldridge
The Minister for Health and Aged Care

This page intentionally left blank

Executive summary

Introduction

The information in this atlas adds to a convincing body of evidence built up over a number of years in Australia as to the striking disparities in health that exist between groups in the population. People of low socioeconomic status (those who are relatively socially or economically deprived) experience worse health than those of higher socioeconomic status for almost every major cause of mortality and morbidity. The challenge for policy makers, health practitioners and governments is to find ways to address these health inequities.

Background

The primary aims of the first edition of *A Social Health Atlas of Australia* were to illustrate the spatial distribution of the socioeconomically disadvantaged population, and to compare this with patterns of distribution of major causes of illness and death and use of health services. The maps and correlation analysis highlighted associations between social and economic factors in relation to health and illness.

A number of new variables have been included in this second edition, together with many of the variables from the first edition. One of the additions is the presentation of data by the new Accessibility/Remoteness Index of Australia (ARIA). Also included is a cluster analysis, providing profiles at the Statistical Local Area (SLA) level of the socioeconomic status, health status and health service utilisation of the population.

The extent of change (between the editions) in the patterns of distribution in death rates by socioeconomic status is also highlighted.

Findings

Correlation analysis

There were correlations of significance at the SLA level between the measures of socioeconomic disadvantage and a number of the health status variables in **Adelaide**. The strongest of these were with the variables for people reporting their health as fair or poor (as opposed to those reporting their health as being excellent, very good, or good); the Physical Component Summary (PCS, a measure of physical health); the handicap status of the population; and premature death from, in particular, circulatory system diseases (**Table 8.1**). Similarly strong associations were evident in the correlation analysis with the variables for the use of GP services by males and females; and of admissions hospital for circulatory and respiratory system diseases, and admissions to a public hospital.

There were fewer correlations of significance at the SLA level in the non-metropolitan areas of South Australia than was the case in **Adelaide**. This is, in part, a result of the number of areas with relatively small numbers of cases (population, deaths, hospital admissions, etc.) which reduces the strength of the analysis. However a number of variables are highly correlated with each other: these are the variables for unemployed people, indigenous Australians, single parent families, people born in non-English speaking countries, people with poor proficiency in English and dwellings without a motor vehicle.

Various sub-sets of these are correlated with measures of health status and use of health services. The strongest correlations with the measures of socioeconomic disadvantage were with the variables for people reporting their health as fair or poor, the PCS and people with a handicap.

Although generally weaker, there was a consistent association between socioeconomic disadvantage and the variables for hospital admissions of males and females; and hospital admissions from circulatory and respiratory system diseases.

Changes in socioeconomic status

Marked variations were recorded between 1986 and 1996 for a majority of the socioeconomic status variables mapped for South Australia (**Table 9.1**). For **Adelaide**, the largest increases were for the population of Aboriginal and Torres Strait Islander people (an increase of 61.2 per cent over ten years); single parent families (36.3 per cent); low income families (32.3 per cent); the occupational grouping of managers and administrators, and professionals (30.0 per cent); people aged 65 years and over (25.8 per cent); unemployed people (18.2 per cent); people born overseas in predominantly non-English speaking countries and resident in Australia for five years or more (12.7 per cent); and dwellings with no motor vehicle (11.3 per cent). The largest decreases recorded over this ten year period were for the variables for unskilled and semi-skilled workers (down by 15.5 per cent) and early school leavers (down by 15.4 per cent).

Variations of this order were also recorded in the non-metropolitan areas of South Australia. The major differences from the changes noted for **Adelaide** were the larger increases in the population of people aged 65 years and over and the number of dwellings without a motor vehicle; smaller increases for indigenous people, single parent families, low income families, unemployed people; and decreases for the occupations of managers and administrators, and professionals, each of the three variables for people born overseas in predominantly non-English speaking countries, and housing authority rented dwellings.

Substantial variations were recorded in income support payments to residents of **Adelaide** for all of the payment types analysed, other than the Age Pension, for which there was a small decrease (a decrease of 1.1 per cent). The number of recipients for each of the other payment types increased substantially, with the number of unemployment beneficiaries almost doubling (an increase of 87.2 per cent) (**Table 9.1**). Similar increases were recorded in the non-metropolitan areas of South Australia for all of these income support payments other than the Age Pension, for which there was a much smaller increase (an increase of 5.1 per cent).

Changes in death rates

Death rates in South Australia have declined over the years 1985 to 1989 and 1992 to 1995 for the majority of causes studied.

In **Adelaide**, the largest decreases were recorded for the infant death rate (down by 30.7 per cent) and for deaths of people aged from 15 to 64 years from circulatory system diseases (down by 36.7 per cent), respiratory system diseases (down by 17.8 per cent) and accidents, poisonings and violence (down by 13.1 per cent). All cause mortality was 17.6 per cent lower over this period, marginally more so for males than for females.

There were reductions in the rates of premature death in the non-metropolitan areas of South Australia for all but cancer (for which there was a slight increase). However the overall reduction was lower than that recorded for **Adelaide**, at around two thirds (69.9 per cent) for all cause mortality.

Summary of findings by socioeconomic status of area of residence

Comparisons are made of changes in the health status of the population by socioeconomic status. In the absence of any direct measure of socioeconomic status in the health status data, the socioeconomic status of the SLA of usual residence in the health status records is used. In this analysis socioeconomic status is measured by the Index of Relative Socio-Economic Status (IRSD, see page 17). The SLAs in **Adelaide** have been grouped into five groups (quintiles) based on the IRSD score, with Quintile 1 comprising the twenty per cent of SLAs with the highest IRSD scores, and Quintile 5 comprising the twenty per cent of SLAs with the lowest IRSD scores. This exercise was repeated for the non-metropolitan areas in South Australia.

Health status

Although there is some variability across the quintiles, the pattern is always for the highest socioeconomic status SLAs (those in Quintile 1) to have the most advantageous (ie. in the majority of cases the lowest) rates and, generally, for the most disadvantaged SLAs (those in Quintile 5) to have the highest rates. The exception is the Physical Component Summary (PCS), for which low scores indicate poorer health (**Figure 9.2**).

Years of potential life lost (YPLL) from deaths between the ages of 15 to 64 years varied from a standardised ratio (SR) in the most advantaged areas of 81 (19 per cent fewer YPLL than were expected from the South Australian State rates) to an SR of 133 in the most disadvantaged areas (indicating that there were 33 per cent more YPLL than were expected from the State rates). Large differentials were also evident for deaths of 15 to 64 year old males (from an SDR of 79 in Quintile to 138 in Quintile 5) and deaths of 15 to 64 years olds from lung cancer (69 to 178), circulatory system diseases (73 to 133) and respiratory system diseases (64 to 138).

The main differences from the gradients evident for **Adelaide** are for deaths of people aged from 15 to 24 years from the combined causes of accidents, poisonings and violence for which, despite the highly elevated ratios, there is no clear gradient.

Health service utilisation

Although there is some variability across the quintiles for the health service utilisation variables for SLAs in **Adelaide**, the pattern is generally for the most advantaged SLAs (those in Quintile 1) to have the lowest rates of admission, and for the most disadvantaged SLAs (those in Quintile 5) to have the highest rates. The exceptions include the graphs for admissions to a private hospital, for all cancers and for breast cancer of females aged 40 years and over, and for the surgical procedures of myringotomy, hip replacement, lens insertion and endoscopy. Others, including the graphs for admissions for psychosis and same day admissions for a surgical procedure, reveal a less consistent pattern (**Figure 9.3**).

The main differences from the gradients evident for **Adelaide** are for admissions for cancer, breast cancer, psychosis, Caesarean section, hip replacement and endoscopy, for which the ratios vary across the quintiles (**Figure 9.5**).

Change in health status by socioeconomic status of area of residence

As noted above, there has been an overall decrease in death rates in South Australia; there are also differentials in death rates by socioeconomic status of area. It is possible to examine the extent of the change in death rates by socioeconomic status of area. As data was not available for non-metropolitan SLAs in the first edition of the atlas, the following comparisons have been limited to **Adelaide**.

Death rates in **Adelaide** declined between 1985-89 and 1992-95 for all of the causes of death studied (with the exception of the 'other' causes group), both overall and in the majority of quintiles of socioeconomic status of area.

It is clear, however, that despite the overall decline, the strong gradient in death rates between the quintiles remains. In fact, the differential in death rates for male residents aged from 15 to 64 years between Quintile 1 (the most advantaged areas) and Quintile 5 (the most disadvantaged areas) increased, from 1.58 times higher in the most disadvantaged areas in 1985-89 to 2.02 times higher in 1992-95. Similar differentials occur for other deaths variables studied.

For females, overall death rates decreased to a similar extent to those for males, and the differential in death rates for female residents aged from 15 to 64 years between Quintile 1 and Quintile 5 also increased, from 1.44 times higher in the most disadvantaged areas in 1985-89 to 1.54 times higher in 1992-95.

Infant death rates in **Adelaide** declined by almost one third (30.7 per cent, **Table 9.2**) between 1985-89 and 1992-95, and the differential in rates between Quintile 1 and Quintile 5 also declined, from 2.34 times higher in the most disadvantaged areas in 1985-89 to 1.64 times higher in 1992-95. Despite this notable reduction, the remaining differential of 64 per cent is still substantial.

Despite a decline in death rates of the 15 to 64 year old population for all cancers and lung cancer (with a larger decline), the differential in rates between Quintile 1 and Quintile 5 increased, from 1.17 times higher in the most disadvantaged areas in 1985-89 to 1.41 times higher in 1992-95 for cancer, and, more substantially so, for lung cancer, from 1.52 to 2.90.

The overall decline in death rates for deaths of 15 to 64 year olds from circulatory system diseases was the highest among the causes of death studied, at over one third (36.7 per cent) in **Adelaide**. The differential in rates between Quintile 1 and Quintile 5 increased from 1.60 times higher in the most disadvantaged areas in 1985-89 to 2.10 times higher in 1992-95.

Although the gradients in deaths rates from respiratory system diseases across the quintiles of socioeconomic status of area of residence in **Adelaide** are not particularly strong, the differential in death rates between Quintiles 1 and 5 is clearly evident. In 1985-89 it was 2.08; by 1992-95 this had increased (by 41.2 per cent) to 2.94. This was the largest differential in death rates for any of the causes studied.

Death rates of 15 to 64 year old people from the external causes of accidents, poisonings and violence are also highest in the most disadvantaged areas of **Adelaide**. Again, the differential in 1992-95 is higher than in 1985-89 (up from 1.59 to 1.74).

As is the case for deaths from the external causes of accidents, poisonings and violence in the 15 to 64 year age group, the differential in death rates for the 15 to 24 year age group in 1992-95 is higher than in 1985-89 (up by 50.4 per cent, from 1.36 to 2.04). The largest declines in death rates were in Quintiles 1 and 3 (down by more than 40 per cent); the smallest of the declines was in Quintile 5 (-8.4 per cent).

Conclusion

There is clear evidence in the data of an association at the SLA level between high premature death rates (both for deaths from all causes and from most specific causes) and socioeconomic disadvantage, as measured by the IRSD. These associations are generally evident not only between the most advantaged (Quintile 1) and disadvantaged areas (Quintile 5), but also at each of the intervening levels of socioeconomic status (Quintiles 2 to 4) (**Figures 9.2 and 9.4**).

Similarly, there are associations between socioeconomic disadvantage and high rates of use of general medical practitioner services in **Adelaide**, and for most of the variables for hospital admission in both **Adelaide** and the non-metropolitan areas of South Australia (**Figures 9.3 and 9.5**). The gradients for admissions by socioeconomic status of area are particularly strong in the non-metropolitan SLAs.

It is also clear that, despite an overall improvement in death rates from all causes and for all of the specific causes studied (with the exception of the 'other causes' group) for **Adelaide** (**Table 9.2, Figure 9.6**), these improvements have not resulted in any significant reduction in the disparities evident in death rates between residents of the most well off areas and those in the poorest areas. In fact, for all but infant deaths, the gap in death rates has increased (**Figure 9.6**).

This page intentionally left blank

Using the *Social Health Atlas*

The social health atlas package

This second edition of *A Social Health Atlas of Australia* comprises:

- this volume for South Australia and a companion volume (Volume 5.2) containing the data mapped (the numbers and rate/ratio/percentages on which the maps are based); and
- similar volumes for each of the other States and Territories and a separate atlas for Australia as a whole (each of these atlases also has a companion volume containing the data mapped).

Some of the data from the atlas are also available on the **HealthWIZ** statistics database product, which comprises comprehensive health statistics from Australia's hospital systems, cause of death registries, population censuses, cancer registries, Medicare and income support system, as well as details of aged care and child care.

This volume contains general background information to the atlas, as well as maps of selected variables showing patterns of socioeconomic status, health status and health and welfare service use at a small area level. Each of these maps is accompanied by a commentary.

The text and maps can also be downloaded for reading and printing from the Public Health Information Development Unit World Wide Web site at www.publichealth.gov.au. The text (including the maps and graphs) and datasets on which the maps are based are available on CD-ROM (for Windows). Further details are in Appendix 1.1, *Project Resources and Output*.

Content

The atlas has nine chapters, an appendix, a bibliography and an index. The chapters are:

- 1 Introduction
- 2 Methods
- 3 Demography and socioeconomic status
- 4 Income support payments
- 5 Health status
- 6 Utilisation of health services
- 7 Availability of selected health services
- 8 Statistical analysis
- 9 Summary

Chapters 1 and 2 provide an overview of the atlas and the approach taken in analysing and mapping data. These sections contain important information on the limitations of the mapped data. The Appendix provides additional background information, and the *Glossary*, at the end of this section, defines some of the terms used.

Chapters 3 to 7 each provide an introduction to the topic(s) being mapped, as well as the maps and associated commentary.

Chapter 8 shows the results of the correlation and cluster analyses. Chapter 9 presents details of the major changes in the data between this second and the first edition, as well as some summary measures of the health differentials calculated from the health status and health service utilisation data mapped in Chapters 5 and 6.

Using the atlas

Some people will use the atlas as a reference source, either going to particular maps (eg. of hospital surgical procedures), or using the index to find a particular topic (eg. deaths from circulatory system diseases) or variable (eg. tonsillectomy).

Others may choose to examine the correlation matrices and to then view the maps for variables for which the data are highly correlated. Or they may access the data in a spreadsheet and re-group the SLAs to suit their own purpose, recalculating the percentages or standardised ratios to represent the new spatial groupings.

To assist users in reading the maps, the layout of the two map types used most frequently is described below. The more detailed discussion in Chapter 2 on the way in which the data have been analysed and presented is, however, important in terms of gaining an understanding of how best to use the data and maps in this atlas. Users of the atlas are particularly encouraged to read this chapter to ensure they are aware of the deficiencies in the datasets presented, as well as in the mapping approach used.

Map of Adelaide

Area mapped

The area mapped is the Statistical Division of **Adelaide** (generally known as the capital city area). The spatial unit mapped is the Statistical Local Area (SLA).

Additional details, including key maps to assist in the location and identification of particular SLAs, are in *Appendix 1.2*: a set of clear film overlays to assist in this process is included in a pocket inside the back cover of this atlas.

Data measures mapped

The map sub-title indicates the format in which the data are presented. In a majority of cases, data are mapped as either a percentage or age (or age-sex) standardised ratio (the process of standardisation is described in Appendix 1.3, *Analysis and presentation of data*). The exceptions are the maps, in Chapter 7, of the location of selected health services; the Index of Relative Socio-Economic Disadvantage mapped in Chapter 3; the infant death rate; and the Total Fertility Rate.

The legend shows the data ranges used to indicate the spatial distribution of the characteristic being mapped.

Footnotes on the map page draw attention to particular aspects of the mapped data and the source of the data.

Description

The text associated with the maps provides background information on the variable being mapped and describes the pattern of distribution of the variable at the SLA level.

The commentary in the top section provides information about the topic being mapped, as well as a comparison between the capital cities and, where the data are available, refers to the situation reported in the first edition of the atlas. For variables where the data are age (or age-sex) standardised, these comparisons are made across Australia (with Australia as the standard for comparison).

In the lower two thirds of the page, attention is drawn to other sources of information about the variable, or characteristics of the population under discussion. The pattern of distribution shown in the map is then described, and associations evident in the correlation analysis with other variables are noted. Users should note that in these descriptions, where data has been standardised, it has been re-calculated to a new standard – in this atlas, to the South Australian State rates (rather than the Australian rates). This allows comparisons to be made between the rates for the SLAs within **Adelaide**, and the South Australian rates – ie. in effect the State average. This differs from the commentary on the top of the page, for which comparisons are made with the Australian rates.

Where the numbers of cases are relatively small (and, in particular, where these small numbers are associated with elevated rates), the absolute numbers are included in the commentary. The numbers (as well as the percentages, rates and ratios) are available in printed and electronic forms and should be used in conjunction with the information in this atlas.

Map of South Australia: referred to as the 'non-metropolitan areas' of South Australia

Area mapped

The spatial units mapped are again SLAs: however **Adelaide** is mapped as one area (ie. not by SLA) to enhance comparisons between the capital city and the non-metropolitan areas.

Towns with a population of 7,500 or more (but less than the urban centre cut-off of 100,000) are represented on the maps as circles. Unfortunately, data for many towns is not available for the datasets in the atlas (other than the Census data).

As noted above in relation to the map of **Adelaide**, additional details are in *Appendix 1.2*: a set of clear film overlays to assist in the location and identification of particular SLAs is included in a pocket inside the back cover of this atlas.

Data measures mapped

See comments above concerning **Adelaide**.

Description

Again, commentary in the top section provides information about the topic being mapped, as well as national comparisons, this time comparing the 'other' major urban centres (those population centres of 100,000 or larger which are not capital cities) and the areas of Australia outside of the capital cities and other major urban centres. These regional/rural/remote areas are referred to in the text as 'non-metropolitan areas'. Where the data are age (or age-sex) standardised, the standard for comparison is, again, Australia.

The lower two thirds of the page again draws attention to other sources of information about the variable, or characteristics of the population under discussion. The pattern of distribution shown in the map is then described, and associations evident in the correlation analysis with other variables are noted. Users should note that in these descriptions, where data has been standardised, it has been re-calculated to a new standard – in this atlas, to the South Australian State rates (rather than the Australian rates). This allows comparisons to be made between the rates for SLAs within the non-metropolitan areas of South Australia and the State rates – ie. in effect the State average.

The cautions in the main introduction and in the introductory notes to each chapter are particularly relevant to the non-metropolitan areas, with their geographically large SLAs and relatively small, scattered populations.

Additional information: ARIA Index

In addition to the map, the map page includes a graph showing the average measure for the variable in each of five levels of accessibility/remoteness, as determined by the Accessibility/Remoteness Index for Australia (ARIA). This Index is described in more detail in Chapter 2, under the heading *Accessibility and Remoteness*. In brief, each SLA in South Australia has been allocated to one of five categories, which range from Highly Accessible, through Accessible, Moderately Accessible and Remote, to Very Remote. The average percentage, rate or ratio for each of the five categories is then calculated for each variable and presented as a graph. The graph is accompanied by a brief comment on the distribution across the categories.

Contents

Chapter	Page
Foreword	iii
Executive summary	v
Using the <i>Social Health Atlas</i>	ix
List of maps	xiv
List of tables	xviii
List of figures	xxii
Acknowledgements	xxiii
Glossary and explanatory notes	xxv
1 Introduction	1
2 Methods	7
3 Demography and socioeconomic status	15
Introduction, data sources and explanatory notes	
Age distribution	
children aged 0 to 4 years	20
people aged 65 years and over	24
Families	
single parent families	28
low income families	32
Labour force	
unskilled and semi-skilled workers	36
unemployed people	38
female labour force participation	42
Educational participation and achievement	
people who left school at age 15 years or less, or did not go to school	46
Aboriginal and Torres Strait Islander people	50
People born in predominantly non-English speaking countries	
number resident in Australia for five years or more	54
number resident in Australia for less than five years	58
proficiency in English	62
Housing	
dwellings rented from State/Territory housing authority	66
dwellings with no motor vehicle	70
SEIFA Index of Relative Socio-economic Disadvantage	74
4 Income support payments	79
Introduction, data sources and explanatory notes	
Age pensioners	82
Disability support pensioners	86
Female sole parent pensioners	90
People receiving an unemployment benefit	94
Dependent children of selected pensioners and beneficiaries	98

Chapter	Page
5 Health status	103
Introduction, data sources and explanatory notes	
Synthetic Predictions of selected health status measures	109
Introduction, data sources and explanatory notes	
People reporting their health as fair or poor	112
Physical Component Summary, SF36	116
Handicap status	120
Deaths	125
Introduction, data sources and explanatory notes	
under one year of age : infant deaths	132
15 to 64 year olds	
all causes: males	134
all causes: females	138
all cancers	142
lung cancer	146
circulatory system diseases	148
respiratory system diseases	152
accidents, poisonings and violence	155
15 to 24 year olds	
accidents, poisonings and violence	162
Years of potential life lost	164
Total Fertility Rate	173
6 Utilisation of health services	179
Introduction, data sources and explanatory notes	
Hospital admissions (including for surgical procedures)	181
Introduction and explanatory notes	
public acute hospitals and private hospitals	192
public acute hospitals	196
private hospitals	200
public acute and private hospitals	
males	204
females	208
same day patients	212
infectious and parasitic diseases	216
all cancers	220
lung cancer	224
cancer of the female breast	226
psychosis	228
neurotic, personality or other mental disorders	232
all circulatory system diseases	236
ischaemic heart disease	240
all respiratory system diseases	244
0 to 4 years olds with respiratory system disease	248
bronchitis, emphysema and asthma	252
accidents, poisonings and violence	256

Chapter	Page
Hospital admissions for surgical procedures	261
Introduction, data sources and explanatory notes	
admissions for a surgical procedure	264
same day admissions for a surgical procedure	268
tonsillectomy and/or adenoidectomy	272
myringotomy	276
Caesarean section	280
hysterectomy	284
hip replacement	288
lens insertion	290
endoscopy	294
General medical practitioner (GP) services	303
Introduction, data sources and explanatory notes	
GP services	
males	306
females	310
Immunisation status of one year old children	314
7 Availability of selected health services	319
Introduction, data sources and explanatory notes	
Population per GP	322
Hospital beds	
public acute hospitals	326
private hospitals	330
Residential care places	
Nursing home places	334
Hostel places	338
8 Statistical analysis	343
Introduction and explanatory notes	
Correlation analysis	343
Cluster analysis	349
9 Summary of findings	367
Introduction	
Changes in data rates between editions	367
Summary of findings by socioeconomic status of area of residence	368
Appendix 1: Supporting documentation	379
1.1 Project resources and output	381
1.2 Geographic areas mapped	383
1.3 Analysis and presentation of data	389
1.4 Classification of deaths, admissions and procedures	391
1.5 Synthetic estimates for small areas	393
1.6 Additional details of cluster analysis	397
Bibliography	403
Index	413

List of maps

Chapter & Map	Page
2 Methods	
2.1 Accessibility/Remoteness Index of Australia (ARIA), 1996	9
2.2 Accessibility/Remoteness Index of Australia (ARIA), for SLAs in South Australia, 1996	10
3 Demography and socioeconomic status	
3.1 Children aged 0 to 4 years, Adelaide, 1996	21
3.2 Children aged 0 to 4 years, South Australia, 1996	23
3.3 People aged 65 years and over, Adelaide, 1996	25
3.4 People aged 65 years and over, South Australia, 1996	27
3.5 Single parent families, Adelaide, 1996	29
3.6 Single parent families, South Australia, 1996	31
3.7 Low income families, Adelaide, 1996	33
3.8 Low income families, South Australia, 1996	35
3.9 Unskilled and semi-skilled workers, Adelaide, 1996	37
3.10 Unemployed people, Adelaide, 1996	39
3.11 Unemployed people, South Australia, 1996	41
3.12 Female labour force participation, Adelaide, 1996	43
3.13 Female labour force participation, South Australia, 1996	45
3.14 People who left school at age 15 years or less, or did not go to school, Adelaide, 1996	47
3.15 People who left school at age 15 years or less, or did not go to school, South Australia, 1996	49
3.16 Aboriginal and Torres Strait Islander people, Adelaide, 1996	51
3.17 Aboriginal and Torres Strait Islander people, South Australia, 1996	53
3.18 People born in predominately non-English speaking countries and resident in Australia for five years or more, Adelaide, 1996	55
3.19 People born in predominately non-English speaking countries and resident in Australia for five years or more, South Australia, 1996	57
3.20 People born in predominately non-English speaking countries and resident for less than five years, Adelaide, 1996	59
3.21 People born in predominately non-English speaking countries and resident for less than five years, South Australia, 1996	61
3.22 Poor proficiency in English of people aged five years and over and born in predominately non-English speaking countries, Adelaide, 1996	63
3.23 Poor proficiency in English of people aged five years and over and born in predominately non-English speaking countries, South Australia, 1996	65
3.24 Dwellings rented from the State housing authority, Adelaide, 1996	67
3.25 Dwellings rented from the State housing authority, South Australia, 1996	69
3.26 Dwellings with no motor vehicles, Adelaide, 1996	71
3.27 Dwellings with no motor vehicles, South Australia, 1996	73
3.28 SEIFA Index of Relative Socio-Economic Disadvantage, Adelaide, 1996	75
3.29 SEIFA Index of Relative Socio-Economic Disadvantage, South Australia, 1996	77
4 Income support payments	
4.1 Age pensioners, Adelaide, 30 June 1996	83
4.2 Age pensioners, South Australia, 30 June 1996	85
4.3 Disability support pensioners, Adelaide, 30 June 1996	87
4.4 Disability support pensioners, South Australia, 30 June 1996	89
4.5 Female sole parent pensioners, Adelaide, 30 June 1996	91
4.6 Female sole parent pensioners, South Australia, 30 June 1996	93
4.7 People receiving an unemployment benefit, Adelaide, 30 June 1996	95
4.8 People receiving an unemployment benefit, South Australia, 30 June 1996	97
4.9 Dependent children of selected pensioners and beneficiaries, Adelaide, 30 June 1996	99
4.10 Dependent children of selected pensioners and beneficiaries, South Australia, 30 June 1996	101

5	Health status	
5.1	People reporting their health as fair or poor, Adelaide, 1995	113
5.2	People reporting their health as fair or poor, South Australia, 1995	115
5.3	Physical Component Summary, SF-36, Adelaide, 1995	117
5.4	Physical Component Summary, SF-36, South Australia, 1995	119
5.5	Estimated number of people with a handicap, Adelaide, 1993	121
5.6	Estimated number of people with a handicap, South Australia, 1993	123
5.7	Infant deaths, Adelaide, 1992 to 1995	133
5.8	Deaths of males aged 15 to 64 years from all causes, Adelaide, 1992 to 1995	135
5.9	Deaths of males aged 15 to 64 years from all causes, South Australia, 1992 to 1995	137
5.10	Deaths of females aged 15 to 64 years from all causes, Adelaide, 1992 to 1995	139
5.11	Deaths of females aged 15 to 64 years from all causes, South Australia, 1992 to 1995	141
5.12	Deaths of people aged 15 to 64 years from cancer, Adelaide, 1992 to 1995	143
5.13	Deaths of people aged 15 to 64 years from cancer, South Australia, 1992 to 1995	145
5.14	Deaths of people aged 15 to 64 years from lung cancer, Adelaide, 1992 to 1995	147
5.15	Deaths of people aged 15 to 64 years from circulatory system diseases, Adelaide, 1992 to 1995	149
5.16	Deaths of people aged 15 to 64 years from circulatory system diseases, South Australia, 1992 to 1995	151
5.17	Deaths of people aged 15 to 64 years from respiratory system diseases, Adelaide, 1992 to 1995.	153
5.18	Deaths of people aged 15 to 64 years from accidents, poisonings and violence, Adelaide, 1992 to 1995	159
5.19	Deaths of people aged 15 to 64 years from accidents, poisonings and violence, South Australia, 1992 to 1995	161
5.20	Deaths of people aged 15 to 24 years from accidents, poisonings and violence, Adelaide, 1992 to 1995	163
5.21	Deaths of people aged 15 to 64 years; years of potential life lost, Adelaide, 1992 to 1995	165
5.22	Deaths of people aged 15 to 64 years; years of potential life lost, South Australia, 1992 to 1995	167
5.23	Total Fertility Rate, Adelaide, 1992 to 1995	175
5.24	Total Fertility Rate, South Australia, 1992 to 1995	177
6	Utilisation of health services	
6.1	Admissions to public acute hospitals and private hospitals, Adelaide, 1995/96	193
6.2	Admissions to public acute hospitals and private hospitals, South Australia, 1995/96	195
6.3	Admissions to public acute hospitals, Adelaide, 1995/96	197
6.4	Admissions to public acute hospitals, South Australia, 1995/96	199
6.5	Admissions to private hospitals, Adelaide, 1995/96	201
6.6	Admissions to private hospitals, South Australia, 1995/96	203
6.7	Admissions of males, Adelaide, 1995/96	205
6.8	Admissions of males, South Australia, 1995/96	207
6.9	Admissions of females, Adelaide, 1995/96	209
6.10	Admissions of females, South Australia, 1995/96	211
6.11	Same day admissions, Adelaide, 1995/96	213
6.12	Same day admissions, South Australia, 1995/96	215
6.13	Admissions for infectious and parasitic diseases, Adelaide, 1995/96	217
6.14	Admissions for infectious and parasitic diseases, South Australia, 1995/96	219

Chapter & Map	Page
6.15 Admissions for cancer, Adelaide, 1995/96	221
6.16 Admissions for cancer, South Australia, 1995/96	223
6.17 Admissions for lung cancer, Adelaide, 1995/96	225
6.18 Admissions of females aged 40 years and over for breast cancer, Adelaide, 1995/96	227
6.19 Admissions for psychosis, Adelaide, 1995/96	229
6.20 Admissions for psychosis, South Australia, 1995/96	231
6.21 Admissions for neurotic, personality or other mental disorders, Adelaide, 1995/96	233
6.22 Admissions for neurotic, personality or other mental disorders, South Australia, 1995/96	235
6.23 Admissions for circulatory system diseases, Adelaide, 1995/96	237
6.24 Admissions for circulatory system diseases, South Australia, 1995/96	239
6.25 Admissions for ischaemic heart disease, Adelaide, 1995/96	241
6.26 Admissions for ischaemic heart disease, South Australia, 1995/96	243
6.27 Admissions for respiratory system disease, Adelaide, 1995/96	245
6.28 Admissions for respiratory system diseases, South Australia, 1995/96	247
6.29 Admissions of children aged 0 to 4 years for respiratory system diseases, Adelaide, 1995/96	249
6.30 Admissions of children aged 0 to 4 years for respiratory system diseases, South Australia, 1995/96	251
6.31 Admissions for bronchitis, emphysema or asthma, Adelaide, 1995/96	253
6.32 Admissions for bronchitis, emphysema or asthma, South Australia, 1995/96	255
6.33 Admissions from accidents, poisonings and violence, Adelaide, 1995/96	257
6.34 Admissions from accidents, poisonings and violence, South Australia, 1995/96	259
6.35 Admissions for a surgical procedure, Adelaide, 1995/96	265
6.36 Admissions for a surgical procedure, South Australia, 1995/96	267
6.37 Same day admissions for a surgical procedure, Adelaide, 1995/96	269
6.38 Same day admissions for a surgical procedure, South Australia, 1995/96	271
6.39 Admissions for tonsillectomy and/or adenoidectomy, Adelaide, 1995/96	273
6.40 Admissions for tonsillectomy and/or adenoidectomy, South Australia, 1995/96	275
6.41 Admissions of children aged 0 to 9 years for a myringotomy, Adelaide, 1995/96	277
6.42 Admissions of children aged 0 to 9 years for a myringotomy, South Australia, 1995/96	279
6.43 Admissions of females aged 15 to 44 years for Caesarean section, Adelaide, 1995/96	281
6.44 Admissions of females aged 15 to 44 years for Caesarean section, South Australia, 1995/96	283
6.45 Admissions of females aged 30 years and over for hysterectomy, Adelaide, 1995/96	285
6.46 Admissions of females aged 30 years and over for hysterectomy, South Australia, 1995/96	287
6.47 Admissions for hip replacement, Adelaide, 1995/96	289
6.48 Admissions for lens insertion, Adelaide, 1995/96	291
6.49 Admissions for lens insertion, South Australia, 1995/96	293
6.50 Admissions for endoscopies, Adelaide, 1995/96	295
6.51 Admissions for endoscopies, South Australia, 1995/96	297
6.52 General medical practitioner services to males, Adelaide, 1996	307
6.53 General medical practitioner services to males, South Australia, 1996	309
6.54 General medical practitioner services to females, Adelaide, 1996	311
6.55 General medical practitioner services to females, South Australia, 1996	313
6.56 Immunisation status of children at 12 months of age, Adelaide, 1998	315
6.57 Immunisation status of children at 12 months of age, South Australia, 1998	317

Chapter & Map	Page
7 Availability of selected health services	
7.1 Population per general medical practitioner, Adelaide, 1996/97	323
7.2 Population per general medical practitioner, South Australia, 1996/97	325
7.3 Public acute hospital beds per 1,000 population, Adelaide, 1995/96	327
7.4 Public acute hospital beds per 1,000 population, South Australia, 1995/96	329
7.5 Private hospital beds per 1,000 population, Adelaide, 30 June 1997	331
7.6 Private hospital beds per 1,000 population, South Australia, 30 June 1997	333
7.7 Nursing home places per 1,000 population aged 70 years and over, Adelaide, 30 June 1997	335
7.8 Nursing home places per 1,000 population aged 70 years and over, South Australia, 30 June 1997	337
7.9 Hostel places per 1,000 population aged 70 years and over, Adelaide, 30 June 1997	339
7.10 Hostel places per 1,000 population aged 70 years and over, South Australia, 30 June 1997	341
8 Statistical analysis	
8.1 Socioeconomic clusters based on Statistical Local Areas in Adelaide	352
8.2 Health status clusters based on Statistical Local Areas in Adelaide	353
8.3 Health service utilisation clusters based on Statistical Local Areas in Adelaide	354
8.4 Social health clusters based on Statistical Local Areas in Adelaide	355
8.5 Socioeconomic clusters based on Statistical Local Areas in South Australia	360
8.6 Health status clusters based on Statistical Local Areas in South Australia	361
8.7 Health service utilisation clusters based on Statistical Local Areas in South Australia	362
8.8 Social health clusters based on Statistical Local Areas in South Australia	363
Appendix 1	
A1 Key map for Statistical Local Areas in Adelaide, 1996	384
A2 Key map for Statistical Local Areas in the non-metropolitan areas of South Australia, 1996	386

List of tables

Chapter & Table	Page
1 Introduction	
1.1 Correlation coefficients for small areas in Adelaide	2
1.2 Small area data of relevance to the National Health Priority Areas	4
2 Methods	
2.1 Conversion of 1996 deaths data to SLA using the ABS Census-based postcode converter: deaths by age group for selected SLAs, South Australia, 1996	12
3 Demography and socioeconomic status	
3.1 Population and area, South Australia, 1996	15
3.2 Population of indigenous Australians, 1986 to 1996	16
3.3 Details of demographic and socioeconomic variables mapped	18
3.4 Proportion of population aged 0 to 4 years, capital cities	20
3.5 Proportion of population aged 0 to 4 years, State/Territory	22
3.6 Proportion of population aged 65 years and over, capital cities	24
3.7 Structure of population aged 65 years and over, South Australia, 1986 and 1996	24
3.8 Proportion of population aged 65 years and over, State/Territory	26
3.9 Single parent families, capital cities	28
3.10 Single parent families, State/Territory	30
3.11 Housing tenure by family type, Adelaide, 1996	30
3.12 Low income families, capital cities	32
3.13 Low income families, State/Territory	34
3.14 Unskilled and semi-skilled workers, capital cities	36
3.15 Unemployed people, capital cities	38
3.16 Unemployed people, State/Territory	40
3.17 Unemployment rates by age, sex and area, South Australia, 1996	40
3.18 Female labour force participation, capital cities	42
3.19 Female labour force participation, State/Territory	44
3.20 People who left school at age 15 years or less, or did not go to school, capital cities	46
3.21 People who left school at age 15 years or less, or did not go to school, State/Territory	48
3.22 Aboriginal and Torres Strait Islander people, capital cities	50
3.23 Aboriginal and Torres Strait Islander people, State/Territory	52
3.24 People born in predominantly non-English speaking countries and resident in Australia for 5 years or more, capital cities	54
3.25 People born in predominantly non-English speaking countries and resident in Australia for 5 years or more, State/Territory	56
3.26 Selected countries of origin of people born in non-English speaking countries, South Australia, 1996	56
3.27 People born in predominantly non-English speaking countries and resident in Australia for less than 5 years, capital cities	58
3.28 People born in predominantly non-English speaking countries and resident in Australia for less than 5 years, State/Territory	60
3.29 Poor proficiency in English of people aged 5 years and over and born in predominantly non-English speaking countries, capital cities	62
3.30 Poor proficiency in English of people aged 5 years and over and born in predominantly non-English speaking countries, State/Territory	64
3.31 Dwellings rented from the State housing authority, capital cities	66
3.32 Dwellings rented from the State housing authority, State/Territory	68
3.33 Dwellings with no motor vehicle, capital cities	70
3.34 Dwellings with no motor vehicle, State/Territory	72
3.35 SEIFA Index of Relative Socio-Economic Disadvantage, capital cities	74
3.36 SEIFA Index of Relative Socio-Economic Disadvantage, State/Territory	76

4 Income support payments

4.1	Income support payments mapped, 30 June 1996	79
4.2	Age pensioners, capital cities	82
4.3	Age pensioners, State/Territory	84
4.4	Disability support pensioners, capital cities	86
4.5	Disability support pensioners, State/Territory	88
4.6	Female sole parent pensioners, capital cities	90
4.7	Female sole parent pensioners, State/Territory	92
4.8	People receiving an unemployment benefit, capital cities	94
4.9	People receiving an unemployment benefit, State/Territory	96
4.10	Dependent children of selected pensioners and beneficiaries, capital cities	98
4.11	Dependent children of selected pensioners and beneficiaries, State/Territory	100

5 Health Status

5.1	Health status indicators by socioeconomic disadvantage of area and sex, Australia, late 1980s	103
5.2	Rate/ratio of mortality inequality by socioeconomic disadvantage of area, 1985-87 and 1995-97	104
5.3	People reporting their health as fair or poor, capital cities	112
5.4	People reporting their health as fair or poor, State/Territory	114
5.5	Physical Component Summary, capital cities, 1995	116
5.6	Physical Component Summary, State/Territory, 1995	118
5.7	Estimated number of people with a handicap, capital cities	120
5.8	Estimated number of people with a handicap, State/Territory	122
5.9	Deaths by cause and age, South Australia, 1992 to 1995	127
5.10	Deaths by selected cause and area, South Australia, 1992 to 1995	128
5.11	Infant deaths, capital cities	132
5.12	Deaths of males aged 15 to 64 years from all causes, capital cities	134
5.13	Deaths of males aged 15 to 64 years from all causes, State/Territory	136
5.14	Deaths of females aged 15 to 64 years from all causes, capital cities	138
5.15	Deaths of females aged 15 to 64 years from all causes, State/Territory	140
5.16	Deaths of people aged 15 to 64 years from cancer, capital cities	142
5.17	Deaths of people aged 15 to 64 years from cancer, State/Territory	144
5.18	Deaths of people aged 15 to 64 years from lung cancer, capital cities	146
5.19	Deaths of people aged 15 to 64 years from circulatory system diseases, capital cities	148
5.20	Deaths of people aged 15 to 64 years from circulatory system diseases, State/Territory	150
5.21	Deaths of people aged 15 to 64 years from respiratory system diseases, capital cities	152
5.22	Deaths from accidents, poisonings & violence, by cause, South Australia, 1992 to 1995	155
5.23	Deaths from accidents, poisonings and violence, by area of residence, South Australia, 1992 to 1995	155
5.24	Deaths of people aged 15 to 64 years from accidents, poisonings and violence, capital cities	158
5.25	Deaths of people aged 15 to 64 years from accidents, poisonings and violence, State/Territory	160
5.26	Deaths of people aged 15 to 24 years from accidents, poisonings and violence, capital cities	162
5.27	Deaths of people aged 15 to 64 years: years of potential life lost, capital cities, 1992 to 1995	164
5.28	Deaths of people aged 15 to 64 years: years of potential life lost, State/Territory, 1992 to 1995	166
5.29	Infant deaths, State/Territory	169
5.30	Deaths of people aged 15 to 64 years from lung cancer, State/Territory	170
5.31	Deaths of people aged 15 to 64 years from respiratory system diseases, State/Territory	171
5.32	Deaths of people aged 15 to 24 years from accidents, poisonings and violence, State/Territory	172
5.33	Total Fertility Rate, capital cities, 1992 to 1995	174
5.34	Total Fertility Rate, State/Territory, 1992 to 1995	176

6	Utilisation of health services	
6.1	Health service use by socioeconomic disadvantage of area and sex, Australia, late 1980s	179
6.2	Admissions of indigenous Australians to public acute and private hospitals, by cause, Australia, 1996/97	182
6.3	Public acute and private hospital admissions included in the analysis, South Australia, 1995/96	188
6.4	Public acute and private hospital admissions, by type of admission: Comparison between editions	190
6.5	Admissions of residents of South Australia by State/Territory of location of hospital, 1995/96	190
6.6	Admissions to public acute hospitals and private hospitals, capital cities	192
6.7	Admissions to public acute hospitals and private hospitals, State/Territory	194
6.8	Admissions to public acute hospitals, capital cities, 1995/96	196
6.9	Admissions to public acute hospitals, State/Territory, 1995/96	198
6.10	Admissions to private hospitals, capital cities, 1995/96	200
6.11	Admissions to private hospitals, State/Territory, 1995/96	202
6.12	Admissions of males, capital cities	204
6.13	Admissions of males, State/Territory	206
6.14	Admissions of females, capital cities	208
6.15	Admissions of females, State/Territory	210
6.16	Same day admissions, capital cities, 1995/96	212
6.17	Same day admissions, State/Territory, 1995/96	214
6.18	Admissions with a principal diagnosis of infectious and parasitic diseases, capital cities	216
6.19	Admissions with a principal diagnosis of infectious and parasitic diseases, State/Territory	218
6.20	Admissions with a principal diagnosis of cancer, capital cities	220
6.21	Admissions with a principal diagnosis of cancer, State/Territory	222
6.22	Admissions with a principal diagnosis of lung cancer, capital cities	224
6.23	Admissions of females aged 40 years and over with a principal diagnosis of breast cancer, capital cities	226
6.24	Admissions with a principal diagnosis of psychosis, capital cities, 1995/96	228
6.25	Admissions with a principal diagnosis of psychosis, State/Territory, 1995/96	230
6.26	Admissions with a principal diagnosis of neurotic, personality or other mental disorders, capital cities, 1995/96	232
6.27	Admissions with a principal diagnosis of neurotic, personality or other mental disorders, State/Territory, 1995/96	234
6.28	Admissions with a principal diagnosis of circulatory system diseases, capital cities	236
6.29	Admissions with a principal diagnosis of circulatory system diseases, State/Territory	238
6.30	Admissions with a principal diagnosis of ischaemic heart disease, capital cities	240
6.31	Admissions with a principal diagnosis of ischaemic heart disease, State/Territory	242
6.32	Admissions with a principal diagnosis of respiratory system diseases, capital cities	244
6.33	Admissions with a principal diagnosis of respiratory system diseases, State/Territory	246
6.34	Admissions of 0 to 4 year olds with a principal diagnosis of respiratory system diseases, capital cities	248
6.35	Admissions of 0 to 4 year olds with a principal diagnosis of respiratory system diseases, State/Territory	250
6.36	Admissions with a principal diagnosis of bronchitis, emphysema or asthma, capital cities	252
6.37	Admissions with a principal diagnosis of bronchitis, emphysema or asthma, State/Territory	254
6.38	Admissions with an external cause of accidents, poisonings and violence, capital cities	256
6.39	Admissions with an external cause of accidents, poisonings and violence, State/Territory	258
6.40	Admission rates for selected sentinel procedures, public and private hospitals, 1996/1997	261
6.41	Standardised admission ratios for selected surgical procedures	261
6.42	Admissions for a surgical procedure, capital cities, 1995/96	264
6.43	Admissions for a surgical procedure, State/Territory, 1995/96	266
6.44	Same day admissions for a surgical procedure, capital cities, 1995/96	268
6.45	Same day admissions for a surgical procedure, State/Territory, 1995/96	270
6.46	Admissions with a principal procedure of tonsillectomy and/or adenoidectomy, capital cities, 1995/96	272
6.47	Admissions with a principal procedure of tonsillectomy and/or adenoidectomy, State/Territory, 1995/96	274
6.48	Admissions of children aged 0 to 9 years with a principal procedure of myringotomy, capital cities, 1995/96	276
6.49	Admissions of children aged 0 to 9 years with a principal procedure of myringotomy, State/Territory, 1995/96	278
6.50	Admissions of females aged 15 to 44 years with a principal procedure of Caesarean section, capital cities, 1995/96	280
6.51	Admissions of females aged 15 to 44 years with a principal procedure of Caesarean section, State/Territory, 1995/96	282
6.52	Admissions of females aged 30 years and over with a principal procedure of hysterectomy, capital cities, 1995/96	284
6.53	Admissions of females aged 30 years and over with a principal procedure of hysterectomy, State/Territory, 1995/96	286

Chapter & Table	Page
6.54 Admissions with a principal procedure of hip replacement, capital cities, 1995/96	288
6.55 Admissions for lens insertion, capital cities, 1995/96	290
6.56 Admissions for lens insertion, State/Territory, 1995/96	292
6.57 Admissions with a principal procedure of endoscopy, capital cities, 1995/96	294
6.58 Admissions with a principal procedure of endoscopy, State/Territory, 1995/96	296
6.59 Admissions with a principal diagnosis of lung cancer, State/Territory	299
6.60 Admissions of females aged 40 years and over with a principal diagnosis of breast cancer, State/Territory	300
6.61 Admissions with a principal diagnosis of hip replacement, State/Territory	301
6.62 Location of Royal Flying Doctor Service bases and number of services, 1997	304
6.63 General medical practitioner services to males, capital cities	306
6.64 General medical practitioner services to males, State/Territory	308
6.65 General medical practitioner services to females, capital cities	310
6.66 General medical practitioner services to females, State/Territory	312
6.67 Proportion of children who were fully immunised at 12 months of age, capital cities, 1998	314
6.68 Proportion of children who were fully immunised at 12 months of age, capital cities, 1998	316
7 Availability of selected health services	
7.1 Patient days for nursing home type patients in public acute hospitals, by area, States and Territories, 1997/98	320
7.2 Nursing home and hostel places per 1,000 population aged 70 years and over, 1997	320
7.3 Population per general medical practitioner, capital cities	322
7.4 Population per general medical practitioner, State/Territory	324
7.5 Public acute hospital beds per 1,000 population, capital cities	326
7.6 Public acute hospital beds per 1,000 population, State/Territory	328
7.7 Private hospitals beds per 1,000 population, capital cities	330
7.8 Private hospital beds per 1,000 population, State/Territory	332
7.9 Nursing home places per 1,000 population aged 70 years and over, capital cities	334
7.10 Nursing home places per 1,000 population aged 70 years and over, State/Territory	336
7.11 Hostel places per 1,000 population aged 70 years and over, capital cities	338
7.12 Hostel places per 1,000 population aged 70 years and over, State/Territory	340
8 Statistical analysis	
8.1 Correlation matrix for SLAs in Adelaide	345
8.2 Correlation matrix for SLAs in the non-metropolitan areas of South Australia	347
8.3 Variables used in cluster analysis	349
8.4 Composition of SLA clusters in Adelaide	350
8.5 Composition of SLA clusters in the non-metropolitan areas of South Australia	358
8.6 Composition of town clusters in Australia	365
9 Summary of findings	
9.1 Changes in demographic and socioeconomic status variables, by Section of State, South Australia	367
9.2 Changes in health status variables, by Section of State, South Australia	368
Appendix	
A1 SLAs not mapped: Population less than 100	383
A2 Urban centres in South Australia	383
A3 Key to Statistical Local Areas in Adelaide, 1996	385
A4 Key to Statistical Local Areas in non-metropolitan areas of South Australia, 1996	387
A5 Data sources	390
A6 ICD-9 Codes for causes of death mapped in Chapter 5	391
A7 ICD-9 Codes for diagnoses/external causes mapped in Chapter 6	391
A8 ICPM Codes for surgical procedures mapped in Chapter 6	391

List of figures

Chapter & Figure	Page
3 Demography and socioeconomic status	
3.1 SEIFA Index of Relative Socio-Economic Disadvantage, capital cities	74
3.2 SEIFA Index of Relative Socio-Economic Disadvantage, Rest of State/Territory	76
4 Income support payments	
4.1 Age pensioners, South Australia, 1996	80
4.2 Disability support pensioners, South Australia, 1996	80
4.3 Female sole parent pensioners, South Australia, 1996	81
4.4 Unemployment beneficiaries, South Australia, 1996	81
5 Health Status	
5.1 Death rates of people aged from 15 to 64 years, by cause, Australia	126
5.2 Death rates of people aged from 15 to 64 years, by cause, South Australia	126
5.3 Deaths from all causes, by age and sex, South Australia, 1992 to 1995	128
5.4 Deaths from cancer, by age and sex, South Australia, 1992 to 1995	129
5.5 Deaths from circulatory system diseases, by age and sex, South Australia, 1992 to 1995	129
5.6 Deaths from respiratory system diseases, by age and sex, South Australia, 1992 to 1995	129
5.7 Deaths from accidents, poisonings and violence, by age and sex, South Australia, 1992 to 1995	130
5.8 Suicide rates of people aged from 25 to 64 years, Adelaide and Rest of State	131
5.9 Suicide rates of people aged from 15 to 24 years, Adelaide and Rest of State	131
5.10 Total Fertility Rate, Adelaide and Rest of State, 1992 to 1995	173
6 Utilisation of health services	
6.1 Admissions to public acute and private hospitals, by age, South Australia and Australia, 1995/96	182
6.2 Admissions to public acute and private hospitals, by age and sex, South Australia, 1995/96	183
6.3 Admissions to public acute hospitals, by age and sex, South Australia, 1995/96	183
6.4 Admissions to private hospitals, by age and sex, South Australia, 1995/96	184
6.5 Same day admissions, by age and sex, South Australia, 1995/96	184
6.6 Admissions for circulatory system diseases, by age and sex, South Australia, 1995/96	185
6.7 Admissions for respiratory system diseases, by age and sex, South Australia, 1995/96	185
6.8 Admissions from accidents, poisonings and violence, by age and sex, South Australia, 1995/96	186
6.9 Admissions for a surgical procedure, by age and sex, South Australia, 1995/96	186
6.10 Same day admissions for a surgical procedure, by age and sex, South Australia, 1995/96	186
6.11 General medical practitioner services, by age and sex, South Australia, 1996	304
9 Summary of findings	
9.1 Differentials in IRSD scores for SLAs in Adelaide	369
9.2 Health status differentials by quintile of socioeconomic disadvantage of area, Adelaide	370
9.3 Health service utilisation differentials by quintile of socioeconomic disadvantage of area, Adelaide	371
9.4 Health status differentials by quintile of socioeconomic disadvantage of area, South Australia	373
9.5 Health service utilisation differentials by quintile of socioeconomic disadvantage of area, South Australia	374
9.6 Change in health status by quintile of socioeconomic disadvantage of area, Adelaide	377

Acknowledgements

The atlas series was produced with the assistance of a number of people and organisations. This atlas for South Australia was produced by John Glover and Sarah Tennant, with the assistance of the other members of the atlas project staff, Kevin Harris and Vija Watts. Sarah was also responsible for day to day management of the data (for checking and storing it), for producing many of the tables and graphs, and for generally keeping in touch with the whole project. Some of the early drafting of Chapter 3 in this atlas was undertaken by David Forster and Cameron Joyce, and Caroline Bruce was responsible for the majority of the work in checking and validating the Census data that was supplied for all of the atlases. These three people moved on to other jobs in the early stages of this project. Lucy Glover checked the data in the drafts against the source material and edited it as necessary. Nicholas Glover inserted many of the map files (as did Kieran Moors) and set up the graphs in Chapter 9.

Outside the Public Health Information Development Unit (PHIDU), Prometheus Information Pty Ltd was the major contributor to the project. Prometheus is contracted by the Commonwealth Department of Health and Aged Care to develop HealthWIZ, the software which was used to produce the maps in this atlas and the data tables (on which the maps are based) in Volume 5.1. Some of the information was already held by Prometheus, and other information needed to be obtained from various Commonwealth, State and Territory agencies and added to the HealthWIZ database in a way that ensured comparability. This was no small task. Although the HealthWIZ software included a mapping facility, the particular approach to publishing the atlas required that special arrangements be made to output the maps in a suitable format. For example, the maps were exported from HealthWIZ and pasted into frames in a MS Word document. Each of these documents was then inserted into the appropriate page in the atlas. Much of the work was highly complex and technical, and required attention to detail and knowledge of the datasets (in particular in identifying potential problems in the data and following these up to confirm or correct them) and statistical geography over a number of years. The quality of the final result, evident in the published product, is testimony to their efforts. George Preston, a Director of the company, was always willing to assist. His knowledge of health statistics and his statistical expertise were frequently of value in making decisions about alternative approaches to the analysis and interpretation of data. Daryl Akerlind and Alain Remont designed the software enhancements to provide the pullouts and town overlays for the maps. Other major contributors at Prometheus were Jane Gorrie and Jennifer Chorley, Zlatan Dzumhur, Jane Lindsay, Jennie Widdowson, Ayse Idehen and (in the earlier stages of the project) Swandi Candra.

In addition to funding the project, the Commonwealth Department of Health and Aged Care took a keen interest in the ongoing work. A number of people had a key role in this, including Ruth Parslow, Karl Higgins and Frances Byers (in aspects of project establishment and contract management) and Jan Bennett and Brendan Gibson (overall direction of the project). In more recent months, Joy Eshpeter and Renata Rustowski had a major role in seeing the project signed off and in

negotiating release arrangements. The support and encouragement of this group of people (and others in the Department) has been greatly appreciated, as were their comments on the final drafts.

The South Australian Department of Human Services had contract responsibility for the atlas for much of the time over which it was produced. They provided a supportive environment in which the atlas could be produced, and made possible the transition of responsibility for the project to PHIDU in April 1999. The support of many people in the Department, including the Chief Executive, Christine Charles, is gratefully acknowledged.

The Australian Institute of Health and Welfare (AIHW) provided the majority of the hospital inpatient data. The AIHW provided other background descriptive material for this chapter, in addition to the data mapped. The main individuals who assisted were Jenny Hargreaves, Janis Shaw and Paul Halliday. The State and Territory health agencies all provided additional details of hospital admissions not available from the AIHW (of admissions of residents of one State or Territory occurring in another).

Colin Mathers of the AIHW and Theo Voss of the Victorian Department of Human Services readily agreed to the use in the atlas of the results of their recent (unpublished) studies into links between socioeconomic status and health status.

All of the data in Chapter 3, as well as a range of other data used throughout the atlas, were purchased from the Australian Bureau of Statistics (ABS). The staff of the Adelaide office of the ABS handled these requests and were thorough and helpful in assisting us to define the data so that it was as comparable as possible with that published in the first edition of the atlas.

The cluster analysis was a major exercise and was undertaken in a highly professional manner by Graeme Tucker. The ARIA graphs and the graphs in Chapter 9 were exported from a module produced by Andrew McAlindon. This module streamlined the calculation of the many rates, percentages etc. used in these sections of the atlas, as well as the production of the final graphs.

Diana Hetzel and Jeanette Pope provided invaluable support in strengthening the discussion of the socioeconomic determinants of health in Chapter 1. Diana contributed in a number of other ways, in particular by providing much of the referenced background material in the topic introductions throughout the atlas; she also read the final drafts. Tony Woollacott and Fearnley Szuster read a number of earlier drafts and Fearnley also provided many useful comments on the later drafts. Thanks are also due to Julie Johninke who produced the cover design, and to Paul Doherty for the photographic image used on the cover.

The final responsibility for the content and comment remains with me.

John Glover
Project Manager
December 1999

This page intentionally left blank

Glossary and Explanatory notes

Cause of death

Causes of death are classified by the Australian Bureau of Statistics to the Ninth (1975) Revision of the World Health Organisation's International Classification of Diseases (ICD-9) which was adopted for world-wide use from 1979.

The cause of death particulars in this publication relate to the underlying cause of death, which the World Health Organisation has defined as the disease or injury which initiated the train of morbid events leading directly to death. Accidental and violent deaths are classified to the circumstances of the accident or violence which produced the fatal injury. Deaths of infants aged less than one month are classified according to the main condition in the infant which contributed to the death.

Details of the ICD-9 codes applicable to the variables mapped in Chapter 5 are shown in *Appendix 1.4*.

Coding of hospital admissions

Diagnoses and procedures are classified according to the International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM October 1988 Revision). External causes are classified according to ICD-9-CM Supplementary Classification of External Causes of Injury and Poisoning ('E' codes) classification codes.

Details of the codes applicable to the variables mapped in Chapter 6 are shown in *Appendix 1.4*.

Admissions

The technical term describing a completed hospital episode (ie. the discharge, death or transfer of a patient) is a 'separation'.

At the time of admission, the age, sex, address of usual residence and other personal details of the patient are recorded. At the end of the episode, at the time of separation from hospital, details of the episode itself are recorded, including the principal diagnosis (and other diagnoses), principal procedure (and other procedures), and the date, time and method (discharge, transfer or death) of separation. Consequently, hospital inpatient data collections are based on separations. In this atlas the more commonly used term of 'admission' has been used. In an analysis such as this, which excludes long stay patients (other than the few long stay acute patients), there is little difference between the number of admissions and the number of separations in a year. Also, 'admission' is a much more familiar term to many people who will use this atlas.

Standardised ratios

Data on which many of the variables have been mapped has been adjusted to remove differences in the data between areas mapped where those differences result from differences in the age and/or sex profiles of the populations being examined. This standardisation process is described in Appendix 1.3, *Analysis and presentation of data*.

Statistical Local Area

The Statistical Local Area (SLA) is a standard geographic area established by the Australian Bureau of Statistics (ABS) to cover the whole of Australia, for the purposes of geographically coding data. It is, in a majority of cases, equivalent to a legal local government area (LGA). SLAs comprise whole LGAs; part LGAs (where the LGA has been split for planning, administrative or statistical purposes); or are unincorporated areas. In South Australia there were 118 LGAs and 130 SLAs at 1 July 1996 (ABS 1996).

Symbols used

- n.a. not available
- .. not applicable
- nil, or less than half the final digit shown
- C City
- DC District Council
- M Municipality
- RC Rural City

This page intentionally left blank